

# Prevalence of Rheumatic Fever and Rheumatic Heart Disease in School Children of Bharateswari Homes of Bangladesh

AAS Majumder<sup>1</sup>, MS Flora<sup>2</sup>, AKMM Islam<sup>1</sup>, M Shahidullah<sup>3</sup>, MA Zafar<sup>3</sup>, R Khanam<sup>3</sup>,  
MA Rashid<sup>3</sup>, MH Iqbal<sup>3</sup>, S Zareen<sup>3</sup>, JU Ahmed<sup>3</sup>,

<sup>1</sup>National Institute of Cardiovascular Diseases; <sup>2</sup>National Institute of Preventive and Social Medicine; <sup>3</sup>National Centre for Control of Rheumatic Fever & Heart Diseases.

## Abstract:

### Key words:

Acute rheumatic fever,  
Rheumatic heart disease,  
Jones' criteria.

**Background:** Rheumatic fever (RF) and rheumatic heart disease (RHD) continue to affect millions of people around the world, including Bangladesh. Children and adolescents are especially susceptible to this disease. Classical risk factors, i.e. poverty, overcrowding, ignorance and insufficient health care services are responsible for the high incidence and prevalence of these diseases. To assess the prevalence of RF and RHD among children, a school survey was conducted in Bharateswari Homes, in the district of Tangail, Bangladesh.

**Methods:** A total of 947 students were examined. Revised Jones' criteria (1992), and clinical examination were used for the diagnosis of RF and RHD.

**Results:** Four cases of RF/RHD were found giving the prevalence of 4.22/1000. This is lower than the prevalence reported in eighties, but is consistent with those found in nineties.

**Conclusion:** Among the school children, there is a declining trend in the prevalence of RF/RHD.

(*Cardiovasc. j.* 2014; 6(2): 103-106)

## Introduction:

RF is a delayed, non-suppurative sequel to upper respiratory tract infection with groups A beta haemolytic (GABH) streptococci. It predominantly affects children and young adults, causes significant morbidity and occasional mortality, and may lead to irreversible damage to the heart valves, i.e. RHD. In the 21st century, RF and RHD are neglected diseases of marginalized communities, though globally, RHD remains the most common cardiovascular disease in young people aged <25 years.<sup>1</sup> Excluding the developed economies, the global burden of RHD in the 5-14 year old children has been estimated to be 0.8-5.7/1000 with a median of 1.3/1000,<sup>2,3</sup> while the overall incidence of acute RF varies from 5 to 51/100,000 population with a mean of 19/100,000.<sup>4</sup> In a recent systematic review, the greatest burden of RF and RHD was found in sub-Saharan Africa, the lowest in North America, the highest mortality rates in the indigenous populations of Australia (23.8/100,000), and among the countries with World Health Organization (WHO) vital registration data, the highest mortality was found in Mauritius (4.32/

100,000).<sup>5</sup> The exact incidence and prevalence of RF and RHD in Bangladesh are not known. Only a limited number of small-scale hospital, school and community surveys are available. In the 2nd half of last century, RF and RHD constituted a significant proportion of admissions in general hospitals, and a lion's share of cardiovascular admissions.<sup>6-8</sup> Probably the community prevalence of RF and RHD was first reported in 1976, which was 7.5/1000 in general population.<sup>9</sup> Subsequent surveys found lower prevalence.<sup>10-12</sup> High prevalence of RF (43.9/1000) and RHD (5.05/1000) was reported in a school survey in Dhaka City in 1984-85.<sup>13</sup> Over the past 3 decades, like many other parts of the world, the incidence and prevalence of acute RF in Bangladesh decreased; however, RHD continues to be an important public health problem here. The present study was planned to know the more recent prevalence of RF and RHD in school children in Bangladesh.

## Materials and Methods:

The study was conducted in Bharateswari Homes (Figs. 1&2), situated in Mirzapur, Tangail,

about 72 km from Dhaka, the capital city of Bangladesh. It is a residential school with children mostly from middle and lower middle class. In the lower classes, both male and female students are enrolled. A total of 947 school children in the age group 4-18 years were screened during the school time. A team comprising of cardiologists, medical graduates, and health assistants visited the school in 2005. Necessary permission was taken from the school authority beforehand. A protocol was prepared based on revised Jones' criteria, 1992 and clinical examination, and for individual student a printed protocol was filled up by the health assistants. A brief history including history of past medical illness was taken and physical examination including detailed cardiovascular examination was done by the physicians. Dynamic auscultation was performed whenever necessary. Sore throat, joint pain with or without swelling, breathlessness on exertion, haemoptysis and ankle oedema were specially observed. Cases with murmurs of grade 2 or above, pan systolic murmur, diastolic murmur and cases with past history of RF but without murmur were noted as suspected cases and were requested to attend Cardiology Department of National Institute of Cardiovascular Diseases (NICVD), Dhaka, for further evaluation.



**Fig.-1:** Bharateswari Homes in Mirzapur, Tangail.



**Fig.-2:** Bharateswari Homes in Mirzapur, Tangail.

### Results:

Of the 947 children, 916 were girls and 31 were boys, so the male female ratio was 1:30. Majority (53.6%) children were in the age group 12-15 years. The mean age of girls was  $12.7 \pm 2.4$  years ranging from 4 to 18 years, while the mean age of boys was  $9.0 \pm 2.2$  years ranging from 5 to 14 years. (Table I). Chest pain and joint pain were present in equal number of cases i.e. 43 (4.5%), fever and breathlessness were found in 4.2% and 2.1% of students respectively. 7.4% students complained of sore throat at the time of examination or in the recent past. A significant cardiac murmur was found in 8 subjects (prevalence rate 8/1000) (Table II). Out of 947 school children, 4 satisfied the clinical diagnosis of RF/RHD, giving a prevalence of 4.22/1000 in general. All the cases were female, and none were male, so the prevalence of RF/RHD was 4.37% in girls. Majority (2) of the RF/RHD cases were in 12-15 year age group (Table III).

**Table-I**

*Age and sex distribution of school children (N=947).*

Age in year	Boys (n=31)		Girls (n=916)		Total (N=947)		p value
	Number	%	Number	%	Number	%	
4-7	9	29.0	26	2.8	35	3.7	
8-11	18	58.1	292	31.9	310	32.7	
12-15	4	12.9	504	55.0	508	53.6	
16-18	0	0.0	94	10.3	94	9.9	
Mean $\pm$	$9.0 \pm 2.2$		$12.7 \pm 2.4$		$12.6 \pm 2.4$		0.001 <sup>S</sup>
SD	(5-14)		(4-18)		(4-18)		

**Table-II**

*Distribution of symptoms and signs of school children (N=947).*

Symptoms and signs	Number	Percentage
Chest pain	43	4.5
Breathlessness	20	2.1
Joint pain	43	4.5
Sore throat	70	7.4
Fever	40	4.2
Murmur	8	0.8

**Table-III**  
*Age distribution of rheumatic fever and  
 rheumatic heart disease cases.*

Age in year	Number of RF cases	Number of RHD cases
8-11	1	0
12-15	2	0
16-19	1	0

### Discussion:

This study was conducted to determine the prevalence of RF/RHD in school children aged 4-18 years in Bangladesh at the advent of new millennium. In this study, out of 947 students, 4 were diagnosed to have RF/RHD. The diagnosis was made clinically, and strict differentiation among acute RF, recurrence of RF and RHD were not made. The murmur found in 8 cases may actually represent RHD cases; however, other possibilities including congenital heart disease and innocent murmur remain. In 1984-85, in a school survey involving 4349 children aged 4-17 years in Dhaka City, the prevalence of RF and RHD was reported as 43.9/1000 and 5.05/1000 respectively,<sup>13</sup> which is much higher than the value obtained in the present study. The prevalence of acute RF was 8.5/1000 in a concurrent study.<sup>14</sup> In late eighties (1989), the prevalence of RF and RHD was found to be 0.85/1000 and 2.8/1000 respectively among 5011 urban school children.<sup>15</sup> In the study involving 10538 school children of Dhaka City, Begum et al. reported the prevalence of RF 2.37/1000, and RHD 0.189/1000 among school children. In both the studies, echocardiography was used in selected cases.<sup>16</sup> Analyzing these studies, it seems that the prevalence of RF/RHD was <10/1000 among school children of Dhaka City at that time, and the unusually high prevalence found by Banoo et al. may be due to some methodological factors. As a result of socioeconomic development and better preventive and promotive services, there was a definite trend towards steady decline in incidence and prevalence of RF in the country.<sup>17</sup> So, the prevalence of 4.22/1000 found in the present study carried out in the first decade of 21<sup>st</sup> century seems to be reasonable. Traditionally, RHD was diagnosed by auscultation for a heart murmur with a stethoscope in those with a history of acute RF. Subsequently,

conventional and portable echocardiography were introduced in studies concerning RF and RHD, and echocardiography has proven to be more sensitive and specific than auscultation.<sup>18</sup> RHD detected on echocardiography without an associated clinically pathological cardiac murmur is referred to as 'subclinical RHD'.<sup>19</sup> Use of echocardiography leads to detection and inclusion of previously unrecognized cases of subclinical carditis, the latter exists at rates up to 10 times higher than that diagnosed by examination alone.<sup>20-25</sup> Recently in 2012, World Heart Federation (WHF) proposed new criteria for echocardiographic diagnosis of RHD.<sup>26</sup> To the best of our knowledge, no studies were carried out to find out the echocardiography-based prevalence of RF/RHD in Bangladesh. So, the prevalence of RF and RHD estimated so far may not be accurate, and the true prevalence of RHD may be much higher in Bangladesh as well.

The main drawback of the present study is that, the diagnosis of RF and RHD was made predominantly on clinical findings. So, there was chance of under-diagnosis of RHD. On the other hand, murmur found in some cases might actually be due to diseases other than RHD. In the study population, there was female preponderance with male female ratio 1:30, so representation by male sex was poor. This was a single-centre study, so the data obtained may not be suitable for generalization to all school children of the country.

### Conclusion:

RF and RHD are still prevalent in Bangladeshi children. In order to include subclinical carditis more efficiently, echocardiography-based studies should be carried out to determine the current 'true' prevalence of RF and RHD. The information available thereby, would help to combat this public health problem more efficiently in future.

---

### Conflict of Interest - None.

---

### References

1. Remenyi B, Carapetis J, Wyber R, Taubert K, Mayosi BM; World Heart Federation. Position statement of the World Heart Federation on the prevention and control of rheumatic heart disease. *Nat Rev Cardiol* 2013 May; 10(5):284-292.

2. Carapetis JR, Steer AC, Mulholland EK, Weber M. The global burden of group A streptococcal disease. *Lancet Infect Dis* 2005; 5:685-694.
3. Carapetis JR. Rheumatic heart disease in Asia. *Circulation* 2008;118:2748-2753.
4. Tibazarwa KB, Volmink JA, Mayosi BM. Incidence of acute rheumatic fever in the world: a systematic review of population-based studies. *Heart* 2008;94:1534-1540.
5. Jackson SJ, Steer AC, Campbell H. Systematic review: Estimation of global burden of non-suppurative sequelae of upper respiratory tract infection: rheumatic fever and post-streptococcal glomerulonephritis. *Trop Med Int Health* 2011 Jan;16(1):2-11.
6. Haque KMHSS, Yoshitake K, Arzu MS, et al. Current status of rheumatic fever and rheumatic heart disease in Bangladesh. *Chest & Heart Bulletin* 1991 Jan; 15:1-4.
7. Khatoon M. Clinical profile of rheumatic fever in some hospitalized children of Bangladesh. *Bangladesh Med Res Counc Bull* 1985;11:1-5.
8. Haque KMHSS. Prevention of rheumatic heart disease. (Editorial) *Bangladesh Heart Journal* 1991;6:1-2.
9. Malik A. Congenital and acquired heart diseases (a survey of 7062 persons). *Bangladesh Med Res Counc Bull* 1976; II: 115-119.
10. Ahmed J, Zaman M, Hassan MM. Prevalence of rheumatic fever and rheumatic heart disease in rural Bangladesh. *Trop Doct* 2005 Jul;35(3):160-161.
11. Haque KMHSS, Hossain M, Mahmud RS, et al. Epidemiology of rheumatic fever and rheumatic heart disease: Observations in 15798 children of rural and urban areas. In: Faruq QO. (ed.) *Proceedings of First National Scientific Conference on Rheumatic Fever and Rheumatic Heart Disease*, May 4-5, 1992, Dhaka, Bangladesh. Dhaka: Ministry of Health and Family Welfare, Government of the People's Republic of Bangladesh and Japan International Cooperation Agency (JICA), Government of Japan; 1992.
12. Hossain MM, Ahmed J, Shahidullah M, Zaman MM. Prevalence of rheumatic fever and rheumatic heart disease in an urban population of Bangladesh. Dhaka: National Centre for Control of Rheumatic Fever and Heart Diseases.
13. Banoo H, Rahman S, Alam A, Azad AK, Sayeed A, Awwal A. Prevalence of rheumatic fever and rheumatic heart diseases in school children in Dhaka city. *Bangladesh Med Res Counc Bull*. 1987 Dec; 13(2):92-100.
14. Janan FAJ. Prevalence of rheumatic fever in an urban population of Bangladesh. *Unpublished Research Results of BMRC Funded Research Project*. Bangladesh Medical Research Council;1991.
15. Mahmud RS, Hossain M, Mosud, et al. Prevalence of rheumatic fever and rheumatic heart diseases in 5-18 years school children of Dhaka city- A study of 5011 school children. *Chest & Heart Bulletin* 1992;16(1):15-22.
16. Begum UHN, Haque KMHSS, Hossain M, Amanullah M, Zafar A. Prevalence of rheumatic fever and rheumatic heart diseases in the Dhaka city, Bangladesh. *Bangladesh Heart Journal* 1994;9(1):4-8.
17. Zaman MM, Yoshiike N, Rouf MA, Mahmud S, Hassan MM. Declining trend of rheumatic fever observed in Bangladesh, 1991-1997. *Trop Doct* 2001 Jul;31(3):169-171.
18. Roberts KV, Brown AD, Maguire GP, Atkinson DN, Carapetis JR. Utility of auscultatory screening for detecting rheumatic heart disease in high-risk children in Australia's Northern Territory. *Med J Australia* 2013 Aug 5;199(3):196-169.
19. Marijon E, Celermajer DS, Tafflet M, et al. Rheumatic heart disease screening by echocardiography: The inadequacy of world health organization criteria for optimizing the diagnosis of subclinical disease. *Circulation* 2009 Aug 25;120(8):663-668.
20. Sadiq M, Islam K, Abid R, Ahmed J, Zaman MM, Keramat Ali SM. Prevalence of rheumatic heart disease in school children of urban Lahore. *Heart* 2009 Mar;95(5):353-357.
21. Paar JA, Berrios NM, Rose JD, et al. Prevalence of rheumatic heart disease in children and young adults in Nicaragua. *Am J Cardiol* 2010 Jun 15;105(12):1809-1814.
22. Bhaya M, Panwar S, Beniwal R, Panwar RB. High prevalence of rheumatic heart disease detected by echocardiography in school children. *Echocardiography* 2010 Apr;27(4):448-453.
23. Saxena A, Ramakrishnan S, Roy A, et al. Prevalence and outcome of subclinical rheumatic heart disease in India: the RHEUMATIC (Rheumatic Heart Echo Utilisation and Monitoring Actuarial Trends in Indian Children) study. *Heart* 2011 Dec;97(24):2018-2022.
24. Baroux N, Rouchon B, Huon B, Germain A, Meunier JM, D'Ortenzio E. High prevalence of rheumatic heart disease in schoolchildren detected by echocardiography screening in New Caledonia. *J Paediatr Child Health* 2013 Feb;49(2):109-114.
25. Rama Kumari N, BhaskaraRaju I, Patnaik AN, et al. Prevalence of rheumatic and congenital heart disease in school children of Andhra Pradesh, South India. *J Cardiovasc Dis Res* 2013 Mar;4(1):11-4.
26. Reményi B, Wilson N, Steer A, et al. World Heart Federation criteria for echocardiographic diagnosis of rheumatic heart disease—an evidence-based guideline. *Nat Rev Cardiol* 2012 Feb 28;9(5):297-309.